

LAPTH-758/99

Vacuum Polarisation Tensors in Constant Electromagnetic Fields: Part I

Christian Schubert

*Laboratoire d'Annecy-le-Vieux de Physique Théorique LAPTH
Chemin de Bellevue, BP 110
F-74941 Annecy-le-Vieux CEDEX
FRANCE
schubert@lapp.in2p3.fr*

Abstract

The string-inspired technique is used for the calculation of vacuum polarisation tensors in constant electromagnetic fields. In the first part of this series, we give a detailed exposition of the method for the case of the QED one-loop N-photon amplitude in a general constant electromagnetic background field. The two-point cases are calculated explicitly, leading to compact representations for the constant field vacuum polarisation tensors for both scalar and spinor QED.

1 Introduction: QED Processes in Constant Electromagnetic Fields

Processes involving constant electromagnetic fields play a special role in quantum electrodynamics. An obvious physical reason is that in many cases a general field can be treated as a constant one to a good approximation. In QED this is expected to be the case if the variation of the field is small on the scale of the electron Compton wavelength. Mathematically, the constant field is distinguished by being one of the very few known field configurations for which the Dirac equation can be solved exactly, allowing one to obtain results which are nonperturbative in the field strength. An early and well-known example is the Euler – Heisenberg Lagrangian [1, 2, 3], the one-loop QED vacuum amplitude in a constant field,

$$\mathcal{L} = \frac{1}{8\pi^2} \int_0^\infty \frac{ds}{s} e^{-ism^2} e^2 ab \frac{\cos(eas) \cosh(ebs)}{\sin(eas) \sinh(ebs)} \quad (1.1)$$

where $a^2 - b^2 \equiv \mathbf{B}^2 - \mathbf{E}^2$, $ab \equiv \mathbf{E} \cdot \mathbf{B}$. This Lagrangian encodes the information on the low energy limit of the one-loop photon S - matrix in a form which is convenient for the derivation of nonlinear QED effects such as photon – photon scattering and vacuum birefringence [4, 5, 6]. Vacuum birefringence is a subject of actual interest [7] since, due to recent improvements in laser technology, the first measurement of this effect in the laboratory seems now imminent [8].

Schwinger’s ingenious use of Fock’s proper–time method in 1951 [9, 10] allowed him to reproduce this result, as well as the analogous one for scalar QED, with considerably less effort. Shortly later Toll [11] initiated the study of the effect of a background field on the one-loop photon propagator. This subject was then over the years investigated by a number of authors, first for the pure magnetic field [12, 13, 14, 15, 16, 17] and crossed field cases [18, 19]. The vacuum polarisation tensor in a general electromagnetic field was first obtained in [20] and given in more explicit form in [21, 22, 23]. The recent [24] contains another recalculation of this quantity, as well as a detailed analysis of the implications for light propagation.

Another consequence of the presence of a background field is the invalidation of Furry’s theorem; already the three-photon amplitude is non-vanishing

in a constant field. Moreover the modification of the photon dispersion relation through the background field can, depending on the photon polarisations, lead to the opening up of phase space for the photon splitting process $\gamma \rightarrow \gamma + \gamma$. For the case of a magnetic field this process was calculated in the low photon energy limit in [25] and for general photon energies in [26]. The amplitude turns out to be very small for the magnetic field strengths presently attainable in the laboratory. Nevertheless, the photon splitting process is believed to be of relevance for the physics of neutron stars which are known to have magnetic fields approaching, and even surpassing, the “critical” magnetic field strength $B_{\text{crit}} = \frac{m_e^2}{e} = 4.41 \times 10^{13}$ Gauss [27, 28, 29]. It seems also not impossible that, with some further improvements in laser technology, photon splitting may be observable in the laboratory in the near future [30, 31].

For QED calculations in constant external fields it is possible and advantageous to take account of the field already at the level of the Feynman rules, i.e. to absorb it into the free electron propagator. Suitable formalisms have been developed decades ago [32, 33, 21]. However beyond the simplest special cases they lead to exceedingly tedious and cumbersome calculations.

A different and more efficient formalism for such calculations has been developed during the last few years, using the so-called “string-inspired” technique. The idea of using string theory methods as a practical tool for calculations in ordinary quantum field theory was advocated by Bern and Kosower [34]. Their work led to the formulation of new computation rules for QCD amplitudes which made it, for example, feasible to perform a complete calculation of the one-loop five gluon amplitudes [35]. A parallel line of work led to the formulation of analogous computation rules for quantum gravity [36, 37]. The relation between the string-derived rules and ordinary Feynman rules was clarified in [38].

Later it was found that even in abelian gauge theory significant improvements over standard field theory methods can be obtained along these lines [39, 40]. Moreover, in this formalism the inclusion of constant external fields turned out to require only relatively minor modifications [41, 42, 43, 44, 45]. For this reason it has been extensively applied to constant field processes in QED. This includes a recalculation of the photon-splitting amplitude [46] as well as high order calculations of the derivative expansion of the QED effective action [42, 43, 47, 48]. A generalization to multiloop photonic am-

plitudes [49, 40, 50, 51, 45] was applied to a calculation of the two-loop correction to the Euler-Heisenberg Lagrangian, using both proper-time [45] and dimensional regularisation [52, 53, 54]. We will not discuss here the involved history of this subject but rather refer the interested reader to the review articles [55, 56]. For other work on QED amplitudes similar in spirit to the string-inspired approach see [57, 58, 59, 60, 61, 62].

In the present paper we first give, in chapter 2, a detailed and self-contained exposition of the string-inspired technique for the calculation of one-loop scalar/spinor QED photon amplitudes in vacuum. In chapter 3 we extend this formalism to the inclusion of constant external fields along the lines of [45]. As a technical improvement on [45] we derive a decomposition of the generalised worldline Green's functions $\mathcal{G}_{B,F}$ in terms of the matrices $\mathbb{1}, F, \tilde{F}, F^2$ with coefficients that are functions of the two standard Maxwell invariants. This will allow us to arrive at explicit results in a manifestly Lorentz covariant way. In chapter 4 we apply the formalism to a calculation of the scalar and spinor QED vacuum polarisation tensors in a constant field. Chapter 5 contains our conclusions.

2 The QED One-Loop N -Photon Amplitude in Vacuum

In [39] it was shown that, for the QED case, the full content of the Bern-Kosower rules can be captured using an approach to quantum field theory based on first-quantized particle path integrals ('worldline path integrals').

2.1 Scalar Quantum Electrodynamics

For the case of scalar QED the basic formulas needed go back to Feynman [63]. The one-loop effective action due to a scalar loop for a Maxwell background can (in modern notation) be written as ¹

¹We work initially in the Euclidean with a positive definite metric $g_{\mu\nu} = \text{diag}(++++)$. The Euclidean field strength tensor is defined by $F^{ij} = \varepsilon_{ijk} B_k, i, j = 1, 2, 3, F^{4i} = -iE_i$, its dual by $\tilde{F}^{\mu\nu} = \frac{1}{2}\varepsilon^{\mu\nu\alpha\beta} F_{\alpha\beta}$ with $\varepsilon^{1234} = 1$. The corresponding Minkowski space amplitudes can be obtained by replacing $g_{\mu\nu} \rightarrow \eta_{\mu\nu} = \text{diag}(-+++)$, $k^4 \rightarrow -ik^0, T \rightarrow iS, \varepsilon^{1234} \rightarrow i\varepsilon^{1230}, \varepsilon^{0123} = 1, F^{4i} \rightarrow F^{0i} = E_i, \tilde{F}^{\mu\nu} \rightarrow -i\tilde{F}^{\mu\nu}$.

$$\Gamma_{\text{scal}}[A] = \int_0^\infty \frac{dT}{T} e^{-m^2 T} \int_{x(T)=x(0)} \mathcal{D}x(\tau) e^{-\int_0^T d\tau \left(\frac{1}{4} \dot{x}^2 + ie \dot{x} \cdot A(x(\tau)) \right)} \quad (2.1)$$

Here T denotes the usual proper-time for the loop fermion. For fixed T $\int \mathcal{D}x$ denotes an integral over the space of all closed loops in spacetime with periodicity T .

This path integral can be used for the calculation of the effective action itself as well as for obtaining the corresponding scattering amplitudes. As a first step in any evaluation, one has to take care of the zero mode contained in it. This is done by fixing the average position of the loop, i.e. one writes

$$\begin{aligned} x^\mu(\tau) &= x_0^\mu + y^\mu(\tau) \\ \int \mathcal{D}x &= \int dx_0 \int \mathcal{D}y \end{aligned} \quad (2.2)$$

where

$$x_0^\mu \equiv \frac{1}{T} \int_0^T d\tau x^\mu(\tau) \quad (2.3)$$

The remaining y path integral is, in the ‘string-inspired formalism’, performed using the Wick contraction rule

$$\langle y^\mu(\tau_1) y^\nu(\tau_2) \rangle = -g^{\mu\nu} G_B(\tau_1, \tau_2) \quad (2.4)$$

where

$$G_B(\tau_1, \tau_2) = |\tau_1 - \tau_2| - \frac{(\tau_1 - \tau_2)^2}{T} \quad (2.5)$$

A “dot” always refers to a derivative in the first variable.

The free Gaussian path integral determinants are, in our conventions, given by

$$\int \mathcal{D}y \, e^{-\int_0^T d\tau \, \frac{1}{4}\dot{y}^2} = (4\pi T)^{-\frac{D}{2}} \quad (2.6)$$

Here D denotes the spacetime dimension. Although in this paper we will consider only the four-dimensional case in this factor D must be left variable in anticipation of dimensional regularization.

We can use this path integral for constructing the scalar QED N -photon amplitude as follows [39]. Expanding the ‘interaction exponential’,

$$\exp\left[-\int_0^T d\tau \, ieA_\mu \dot{x}^\mu\right] = \sum_{N=0}^{\infty} \frac{(-ie)^N}{N!} \prod_{i=0}^N \int_0^T d\tau_i \left[\dot{x}^\mu(\tau_i) A_\mu(x(\tau_i))\right] \quad (2.7)$$

the individual terms correspond to Feynman diagrams describing a fixed number of interactions of the scalar loop with the external field. The corresponding N – photon scattering amplitude is then obtained by specializing to a background consisting of a sum of plane waves with definite polarizations,

$$A_\mu(x) = \sum_{i=1}^N \varepsilon_{i\mu} e^{ik_i \cdot x} \quad (2.8)$$

and picking out the term containing every ε_i once. This immediately yields the following representation for the N - photon amplitude,

$$\begin{aligned} \Gamma_{\text{scal}}[k_1, \varepsilon_1; \dots; k_N, \varepsilon_N] &= (-ie)^N \int_0^\infty \frac{dT}{T} e^{-m^2 T} (4\pi T)^{-\frac{D}{2}} \\ &\times \left\langle V_A^0[k_1, \varepsilon_1] \dots V_A^0[k_N, \varepsilon_N] \right\rangle \end{aligned} \quad (2.9)$$

Here V_A^0 denotes the same photon vertex operator which is also used in string perturbation theory (see, e.g., [64]),

$$V_A^0[k, \varepsilon] \equiv \int_0^T d\tau \, \varepsilon \cdot \dot{x}(\tau) e^{ikx(\tau)} \quad (2.10)$$

At this stage the zero-mode integration (2.2) can be performed, yielding the energy-momentum conservation factor

$$\int d^D x_0 \prod_{i=1}^N e^{ik_i \cdot x_0} = (2\pi)^D \delta(\sum k_i) \quad (2.11)$$

The reduced path integral $\int \mathcal{D}y$ is Gaussian. Its evaluation therefore amounts to Wick contracting the expression

$$\left\langle \dot{y}_1^{\mu_1} e^{ik_1 \cdot y_1} \dots \dot{y}_N^{\mu_N} e^{ik_N \cdot y_N} \right\rangle \quad (2.12)$$

using the correlator (2.4). For the performance of the Wick contractions it is convenient to formally exponentiate all the $\dot{y}_i$'s, writing

$$\varepsilon_i \cdot \dot{y}_i e^{ik_i \cdot y_i} = e^{\varepsilon_i \cdot \dot{y}_i + ik_i \cdot y_i} \big|_{\text{lin}(\varepsilon_i)} \quad (2.13)$$

This allows one to rewrite the product of N photon vertex operators as an exponential. Then one needs only to ‘complete the square’ to arrive at the following closed expression for the one-loop N - photon amplitude [39]

$$\begin{aligned} \Gamma_{\text{scal}}[k_1, \varepsilon_1; \dots; k_N, \varepsilon_N] &= (-ie)^N (2\pi)^D \delta(\sum k_i) \\ &\times \int_0^\infty \frac{dT}{T} (4\pi T)^{-\frac{D}{2}} e^{-m^2 T} \prod_{i=1}^N \int_0^T d\tau_i \\ &\times \exp \left\{ \sum_{i,j=1}^N \left[\frac{1}{2} G_{Bij} k_i \cdot k_j + i \dot{G}_{Bij} k_i \cdot \varepsilon_j + \frac{1}{2} \ddot{G}_{Bij} \varepsilon_i \cdot \varepsilon_j \right] \right\} \big|_{\text{multi-linear}} . \end{aligned} \quad (2.14)$$

Here it is understood that only the terms linear in all the $\varepsilon_1, \dots, \varepsilon_N$ have to be taken. Besides the Green's function G_B also its first and second derivatives appear,

$$\begin{aligned} \dot{G}_B(\tau_1, \tau_2) &= \text{sign}(\tau_1 - \tau_2) - 2 \frac{(\tau_1 - \tau_2)}{T} \\ \ddot{G}_B(\tau_1, \tau_2) &= 2\delta(\tau_1 - \tau_2) - \frac{2}{T} \end{aligned} \quad (2.15)$$

With ‘dots’ we generally denote a derivative acting on the first variable, $\dot{G}_B(\tau_1, \tau_2) \equiv \frac{\partial}{\partial \tau_1} G_B(\tau_1, \tau_2)$, and we abbreviate $G_{Bij} \equiv G_B(\tau_i, \tau_j)$ etc.

The expression (2.14) is identical with the corresponding special case of the ‘Bern-Kosower Master Formula’ [34]. Let us consider explicitly the vacuum polarisation case, $N = 2$. For $N = 2$ the expansion of the exponential factor yields the following expression,

$$\left(\ddot{G}_{B12} \varepsilon_1 \cdot \varepsilon_2 + \dot{G}_{B12}^2 \varepsilon_1 \cdot k_2 \varepsilon_2 \cdot k_1 \right) e^{G_{B12} k_1 \cdot k_2} \quad (2.16)$$

After performing a partial integration on the first term of eq. (2.16) in either τ_1 or τ_2 , the integrand turns into

$$\left(\varepsilon_1 \cdot \varepsilon_2 k^2 - \varepsilon_1 \cdot k \varepsilon_2 \cdot k \right) \dot{G}_{B12}^2 e^{-G_{B12} k^2} \quad (2.17)$$

($k = k_1 = -k_2$). Thus we have

$$\begin{aligned} \Gamma_{\text{scal}}[k_1, \varepsilon_1; k_2, \varepsilon_2] &= (2\pi)^D \delta(k_1 + k_2) \varepsilon_1 \cdot \Pi_{\text{scal}}(k) \cdot \varepsilon_2 \\ \Pi_{\text{scal}}^{\mu\nu}(k) &= \frac{e^2}{(4\pi)^{\frac{D}{2}}} (k^\mu k^\nu - g^{\mu\nu} k^2) \int_0^\infty \frac{dT}{T} T^{-\frac{D}{2}} e^{-m^2 T} \\ &\quad \times \prod_{i=1}^2 \int_0^T d\tau_i \dot{G}_{B12}^2 e^{-G_{B12} k^2} \end{aligned} \quad (2.18)$$

Note that the transversality of the vacuum polarization tensor is already manifest. We rescale to the unit circle, $\tau_i = T u_i, i = 1, 2$, and use translation invariance in τ to fix the zero to be at the location of the second vertex operator, $u_2 = 0, u_1 = u$. We have then

$$G_B(\tau_1, \tau_2) = T u(1 - u), \quad \dot{G}_B(\tau_1, \tau_2) = 1 - 2u \quad (2.19)$$

After performing the trivial T - integration one arrives at

$$\begin{aligned} \Pi_{\text{scal}}^{\mu\nu}(k) &= \frac{e^2}{(4\pi)^{\frac{D}{2}}} (k^\mu k^\nu - g^{\mu\nu} k^2) \Gamma\left(2 - \frac{D}{2}\right) \\ &\quad \times \int_0^1 du (1 - 2u)^2 \left[m^2 + u(1 - u) k^2 \right]^{\frac{D}{2} - 2} \end{aligned} \quad (2.20)$$

The result of the final integration is, of course, the same as is found in the standard field theory calculation for the sum of the corresponding two Feynman diagrams (see, e.g., [65]).

2.2 Spinor Quantum Electrodynamics

The worldline path integral representation (2.1) can be generalized to the spinor QED case in various different ways. The formulation most suitable to the ‘stringy’ approach uses Grassmann variables [66, 67, 68],

$$\begin{aligned} \Gamma_{\text{spin}}[A] = & -\frac{1}{2} \int_0^\infty \frac{dT}{T} e^{-m^2 T} \int \mathcal{D}x \int \mathcal{D}\psi \\ & \times \exp \left[- \int_0^T d\tau \left(\frac{1}{4} \dot{x}^2 + \frac{1}{2} \psi \cdot \dot{\psi} + ieA \cdot x - ie\psi \cdot F \cdot \psi \right) \right] \end{aligned} \quad (2.21)$$

Thus we have, in addition to the same coordinate path integral as in (2.1), a Grassmann path integral $\int \mathcal{D}\psi$ representing the fermion spin. The boundary conditions on the Grassmann path integral are antiperiodic, $\psi(T) = -\psi(0)$, so that there is no new zero mode. The appropriate correlator is

$$\langle \psi^\mu(\tau_1) \psi^\nu(\tau_2) \rangle = \frac{1}{2} g^{\mu\nu} G_F(\tau_1, \tau_2) \quad (2.22)$$

Our normalization for the free Grassmann path integral is

$$\int \mathcal{D}\psi e^{-\int_0^T d\tau \frac{1}{2} \psi \cdot \dot{\psi}} = 4 \quad (2.23)$$

The photon vertex operator (2.10) acquires an additional Grassmann piece,

$$V_A^{\frac{1}{2}}[k, \varepsilon] \equiv \int_0^T d\tau \left(\varepsilon \cdot \dot{x} + 2i\varepsilon \cdot \psi k \cdot \psi \right) e^{ikx} \quad (2.24)$$

Looking again at the vacuum polarization case, we need to Wick-contract two copies of the above vertex operator. The calculation of $\int \mathcal{D}x$ is identical with the scalar QED calculation. The additional contribution from $\int \mathcal{D}\psi$ is

$$(2i)^2 \langle \psi_1^\mu \psi_1 \cdot k_1 \psi_2^\nu \psi_2 \cdot k_2 \rangle = -G_{F12}^2 (g^{\mu\nu} k^2 - k^\mu k^\nu) = -(g^{\mu\nu} k^2 - k^\mu k^\nu) \quad (2.25)$$

Taking the free Grassmann path integral normalization (2.23) into account, eq.(2.18) for the scalar QED vacuum polarisation tensor generalises to the spinor QED case as follows,

$$\begin{aligned} \Pi_{\text{spin}}^{\mu\nu}(k) &= -2 \frac{e^2}{(4\pi)^{\frac{D}{2}}} (k^\mu k^\nu - g^{\mu\nu} k^2) \int_0^\infty \frac{dT}{T} T^{-\frac{D}{2}} e^{-m^2 T} \\ &\quad \times \prod_{i=1}^2 \int_0^T d\tau_i (\dot{G}_{B12}^2 - G_{F12}^2) e^{-G_{B12} k^2} \end{aligned} \quad (2.26)$$

Proceeding as before one obtains

$$\begin{aligned} \Pi_{\text{spin}}^{\mu\nu}(k) &= 8 \frac{e^2}{(4\pi)^{\frac{D}{2}}} (k^\mu k^\nu - g^{\mu\nu} k^2) \Gamma(2 - \frac{D}{2}) \\ &\quad \times \int_0^1 du u(1-u) [m^2 + u(1-u)k^2]^{\frac{D}{2}-2} \end{aligned} \quad (2.27)$$

Remarkably, the explicit calculation of the Grassmann path integral can be circumvented, and replaced by the following simple pattern matching rule [34]. Writing out the exponential in eq.(2.14) one obtains an integrand

$$\exp\left\{\cdots\right\} \big|_{\text{multi-linear}} = (-i)^N P_N(\dot{G}_{Bij}, \ddot{G}_{Bij}) \exp\left[\frac{1}{2} \sum_{i,j=1}^N G_{Bij} k_i \cdot k_j\right] \quad (2.28)$$

with a certain polynomial P_N depending on the various $\dot{G}_{Bij}, \ddot{G}_{Bij}$ and on the kinematic invariants. Now one removes all second derivatives $\ddot{G}_{Bij}$ appearing in P_N by suitable partial integrations in the variables τ_i ,

$$P_N(\dot{G}_{Bij}, \ddot{G}_{Bij}) e^{\frac{1}{2} \sum G_{Bij} k_i \cdot k_j} \xrightarrow{\text{part.int.}} Q_N(\dot{G}_{Bij}) e^{\frac{1}{2} \sum G_{Bij} k_i \cdot k_j} \quad (2.29)$$

This is possible for any N [34]. The result is an alternative integrand for the scalar QED amplitude involving only G_B and $\dot{G}_B$. The integrand for the spinor loop case can then, up to the global factor of -2 , be obtained from the one for the scalar loop simply by replacing every closed cycle of $\dot{G}_B$'s appearing in Q_N by its “worldline supersymmetrization”,

$$\dot{G}_{Bi_1i_2}\dot{G}_{Bi_2i_3}\cdots\dot{G}_{Bi_ni_1}\rightarrow\dot{G}_{Bi_1i_2}\dot{G}_{Bi_2i_3}\cdots\dot{G}_{Bi_ni_1}-G_{Fi_1i_2}G_{Fi_2i_3}\cdots G_{Fi_ni_1}\quad (2.30)$$

Note that an expression is considered a cycle already if it can be put into cycle form using the antisymmetry of $\dot{G}_B$ (e.g. $\dot{G}_{B12}\dot{G}_{B12}=-\dot{G}_{B12}\dot{G}_{B21}$). The replacement is done simultaneously on all cycles.

For $N > 3$ the result of the partial integration procedure is not unique, however the above replacement rule is valid for all possible results. In [69] a certain standardized way was found for performing the partial integrations which leads to a canonical, permutation symmetric and gauge invariant decomposition of the QED N - photon amplitudes.

3 The QED N-Photon Amplitude in a Constant Field

3.1 Generalization of the Bern-Kosower Master Formula

The presence of an additional constant external field, taken in Fock-Schwinger gauge centered at x_0 [41], changes the path integral Lagrangian in eq.(2.21) only by a term quadratic in the fields, $\Delta\mathcal{L}=\frac{1}{2}ie y^\mu F_{\mu\nu}\dot{y}^\nu-ie\psi^\mu F_{\mu\nu}\dot{\psi}^\nu$. The field can therefore be absorbed by a change of the free worldline propagators, replacing $G_B, \dot{G}_B, G_F$ by ([42, 44, 45]; see also [62])

$$\mathcal{G}_B(\tau_1, \tau_2) = \frac{T}{2\mathcal{Z}^2} \left(\frac{\mathcal{Z}}{\sin(\mathcal{Z})} e^{-i\mathcal{Z}\dot{G}_{B12}} + i\mathcal{Z}\dot{G}_{B12} - 1 \right) \quad (3.1)$$

$$\dot{\mathcal{G}}_B(\tau_1, \tau_2) = \frac{i}{\mathcal{Z}} \left(\frac{\mathcal{Z}}{\sin(\mathcal{Z})} e^{-i\mathcal{Z}\dot{G}_{B12}} - 1 \right) \quad (3.2)$$

$$\mathcal{G}_F(\tau_1, \tau_2) = G_{F12} \frac{e^{-i\mathcal{Z}\dot{G}_{B12}}}{\cos(\mathcal{Z})} \quad (3.3)$$

where we have defined $\mathcal{Z} \equiv eFT$. These expressions should be understood as power series in the field strength matrix F . Note that the generalized Green's functions are still translationally invariant in τ , and thus functions of $\tau_1 - \tau_2$. By writing them as functions of the vacuum worldline Green's functions $\dot{G}_B, G_F$ we have left the τ - dependence implicit. This allows us to avoid making a case distinction between $\tau_1 > \tau_2$ and $\tau_1 < \tau_2$ that would become necessary otherwise [44]. Note also the symmetry properties

$$\mathcal{G}_B(\tau_1, \tau_2) = \mathcal{G}_B^T(\tau_2, \tau_1), \quad \dot{\mathcal{G}}_B(\tau_1, \tau_2) = -\dot{\mathcal{G}}_B^T(\tau_2, \tau_1), \quad \mathcal{G}_F(\tau_1, \tau_2) = -\mathcal{G}_F^T(\tau_2, \tau_1) \quad (3.4)$$

Since $\mathcal{G}_B, \mathcal{G}_F$ are, in general, nontrivial Lorentz matrices, the Wick contraction rules eqs.(2.4),(2.22) have to be replaced by

$$\langle y^\mu(\tau_1) y^\nu(\tau_2) \rangle = -\mathcal{G}_B^{\mu\nu}(\tau_1, \tau_2) \quad (3.5)$$

$$\langle \psi^\mu(\tau_1) \psi^\nu(\tau_2) \rangle = \frac{1}{2} \mathcal{G}_F^{\mu\nu}(\tau_1, \tau_2) \quad (3.6)$$

Another slight complication compared to the vacuum case is that, in contrast to their vacuum counterparts, $\mathcal{G}_B, \dot{\mathcal{G}}_B$, and $\mathcal{G}_F$ have non-vanishing coincidence limits. Those are τ - independent:

$$\mathcal{G}_B(\tau, \tau) = \frac{T}{2\mathcal{Z}^2} (\mathcal{Z} \cot(\mathcal{Z}) - 1) \quad (3.7)$$

$$\dot{\mathcal{G}}_B(\tau, \tau) = i \cot(\mathcal{Z}) - \frac{i}{\mathcal{Z}} \quad (3.8)$$

$$\mathcal{G}_F(\tau, \tau) = -i \tan(\mathcal{Z}) \quad (3.9)$$

They are obtained from eqs.(3.1),(3.2),(3.3) using the rules that

$$\dot{G}_B(\tau, \tau) = 0, \quad \dot{G}_B^2(\tau, \tau) = 1 \quad (3.10)$$

This is almost all we need to know for computing one-loop photon scattering amplitudes, or the corresponding effective action, in a constant overall background field. The only further information required at the one-loop level

is the change in the free path integral determinants due to the external field. This change is [41]

$$(4\pi T)^{-\frac{D}{2}} \rightarrow (4\pi T)^{-\frac{D}{2}} \det^{-\frac{1}{2}} \left[\frac{\sin(\mathcal{Z})}{\mathcal{Z}} \right] \quad (\text{Scalar QED}) \quad (3.11)$$

$$(4\pi T)^{-\frac{D}{2}} \rightarrow (4\pi T)^{-\frac{D}{2}} \det^{-\frac{1}{2}} \left[\frac{\tan(\mathcal{Z})}{\mathcal{Z}} \right] \quad (\text{Spinor QED}) \quad (3.12)$$

Since those determinants describe the vacuum amplitude in a constant field they turn out to be, of course, just the proper-time integrands of the Euler-Heisenberg-Schwinger formulas (see (1.1)).

Retracing our above calculation of the N - photon path integral with the external field included we arrive at the following generalization of eq.(2.14), representing the scalar QED N - photon scattering amplitude in a constant field [44, 45]:

$$\begin{aligned} \Gamma_{\text{scal}}[k_1, \varepsilon_1; \dots; k_N, \varepsilon_N] &= (-ie)^N (2\pi)^D \delta(\sum k_i) \\ &\times \int_0^\infty \frac{dT}{T} [4\pi T]^{-\frac{D}{2}} e^{-m^2 T} \det^{-\frac{1}{2}} \left[\frac{\sin(\mathcal{Z})}{\mathcal{Z}} \right] \prod_{i=1}^N \int_0^T d\tau_i \\ &\times \exp \left\{ \sum_{i,j=1}^N \left[\frac{1}{2} k_i \cdot \mathcal{G}_{Bij} \cdot k_j - i\varepsilon_i \cdot \dot{\mathcal{G}}_{Bij} \cdot k_j + \frac{1}{2} \varepsilon_i \cdot \ddot{\mathcal{G}}_{Bij} \cdot \varepsilon_j \right] \right\} \Big|_{\text{multi-linear}} \end{aligned} \quad (3.13)$$

From this formula it is obvious that adding a constant matrix to $\mathcal{G}_B$ will have no effect due to momentum conservation. We can use this fact to get rid of the coincidence limit of $\mathcal{G}_B$, (3.7), namely instead of $\mathcal{G}_B$ one can work with the equivalent Green's function $\bar{\mathcal{G}}_B$, defined by

$$\bar{\mathcal{G}}_B(\tau_1, \tau_2) \equiv \mathcal{G}_B(\tau_1, \tau_2) - \mathcal{G}_B(\tau, \tau) \quad (3.14)$$

No such redefinition is possible for $\dot{\mathcal{G}}_B$ or $\mathcal{G}_F$.

The transition from scalar to spinor QED is done as in the vacuum case, again with only some minor modifications. The spinor QED integrand for a given number of photon legs N is obtained from the scalar QED integrand by the following generalization of the Bern-Kosower algorithm:

1. *Partial Integration:* After expanding out the exponential in the master formula (3.13), and taking the part linear in all $\varepsilon_1, \dots, \varepsilon_N$, remove all second derivatives $\ddot{\mathcal{G}}_B$ appearing in the result by suitable partial integrations in $\tau_1, \dots, \tau_N$.
2. *Replacement Rule:* Apply to the resulting new integrand the replacement rule (2.30) with $\dot{G}_B, G_F$ substituted by $\dot{\mathcal{G}}_B, \mathcal{G}_F$. Since the Green's functions $\mathcal{G}_B, \mathcal{G}_F$ are, in contrast to their vacuum counterparts, non-trivial matrices in the Lorentz indices, it must be mentioned here that the cycle property is defined solely in terms of the τ – indices, irrespective of what happens to the Lorentz indices. For example, the expression

$$\varepsilon_1 \cdot \dot{\mathcal{G}}_{B12} \cdot k_2 \varepsilon_2 \cdot \dot{\mathcal{G}}_{B23} \cdot \varepsilon_3 k_3 \cdot \dot{\mathcal{G}}_{B31} \cdot k_1$$

would have to be replaced by

$$\varepsilon_1 \cdot \dot{\mathcal{G}}_{B12} \cdot k_2 \varepsilon_2 \cdot \dot{\mathcal{G}}_{B23} \cdot \varepsilon_3 k_3 \cdot \dot{\mathcal{G}}_{B31} \cdot k_1 - \varepsilon_1 \cdot \mathcal{G}_{F12} \cdot k_2 \varepsilon_2 \cdot \mathcal{G}_{F23} \cdot \varepsilon_3 k_3 \cdot \mathcal{G}_{F31} \cdot k_1$$

The only other difference compared to the vacuum case is due to the non-vanishing coincidence limits of $\dot{\mathcal{G}}_B, \mathcal{G}_F$, eqs.(3.8),(3.9). Those lead to an extension of the “cycle replacement rule” to include one-cycles [45]:

$$\dot{\mathcal{G}}_B(\tau_i, \tau_i) \rightarrow \dot{\mathcal{G}}_B(\tau_i, \tau_i) - \mathcal{G}_F(\tau_i, \tau_i) = \frac{i}{\sin(\mathcal{Z})\cos(\mathcal{Z})} - \frac{i}{\mathcal{Z}} \quad (3.15)$$

3. The scalar QED Euler-Heisenberg-Schwinger determinant factor must be replaced by its spinor QED equivalent,

$$\det^{-\frac{1}{2}} \left[\frac{\sin(\mathcal{Z})}{\mathcal{Z}} \right] \rightarrow \det^{-\frac{1}{2}} \left[\frac{\tan(\mathcal{Z})}{\mathcal{Z}} \right] \quad (3.16)$$

4. Multiply by the usual factor of -2 for statistics and degrees of freedom.

3.2 Lorentz covariant decomposition of the generalized worldline Green's functions

For the result to be practically useful it will be necessary to know $\mathcal{G}_B, \mathcal{G}_F$ in more explicit form. In the calculations performed in [45, 46, 52] for a purely magnetic field $\mathbf{B}$ pointing along the z - direction the explicit matrix form of $\mathcal{G}_B, \mathcal{G}_F$ had been used. This would also be possible in the generic case, where, excepting the case $\mathbf{E} \cdot \mathbf{B} = 0$, one could use the Lorentz invariance to choose both $\mathbf{E}$ and $\mathbf{B}$ to point along the z - axis. For this case the Green's functions can be easily written out explicitly. However, it is possible to directly express them in terms of Lorentz invariants, without specialisation of the Lorentz frame. This can be done in the following way. Defining the Maxwell invariants

$$\begin{aligned} f &\equiv \frac{1}{4}F_{\mu\nu}F_{\mu\nu} = \frac{1}{2}(B^2 - E^2) \\ g &\equiv \frac{1}{4}F_{\mu\nu}\tilde{F}_{\mu\nu} = i\mathbf{E} \cdot \mathbf{B} \end{aligned} \tag{3.17}$$

we have the relations

$$F^2 + \tilde{F}^2 = -2f\mathbb{1} \tag{3.18}$$

$$F\tilde{F} = -g\mathbb{1} \tag{3.19}$$

Define

$$F_{\pm} \equiv \frac{N_{\pm}^2 F - N_+ N_- \tilde{F}}{N_{\pm}^2 - N_{\mp}^2} \tag{3.20}$$

$$N_{\pm} \equiv n_+ \pm n_- \tag{3.21}$$

$$n_{\pm} \equiv \sqrt{\frac{f \pm g}{2}} \tag{3.22}$$

Then one has

$$F = F_+ + F_- \quad (3.23)$$

$$F^2 F_{\pm} = -N_{\pm}^2 F_{\pm} \quad (3.24)$$

$$F_+ F_- = 0 \quad (3.25)$$

With the help of these relations one easily derives the following formulas,

$$\begin{aligned} f_{\text{even}}(F) &= \frac{1}{N_+^2 - N_-^2} \left\{ -f_{\text{even}}(iN_+) [N_-^2 \mathbb{1} + F^2] + f_{\text{even}}(iN_-) [N_+^2 \mathbb{1} + F^2] \right\} \\ f_{\text{odd}}(F) &= \frac{i}{N_+^2 - N_-^2} \left\{ [N_- f_{\text{odd}}(iN_-) - N_+ f_{\text{odd}}(iN_+)] F \right. \\ &\quad \left. + [N_- f_{\text{odd}}(iN_+) - N_+ f_{\text{odd}}(iN_-)] \tilde{F} \right\} \end{aligned} \quad (3.26)$$

where f_{even} (f_{odd}) are arbitrary even (odd) functions in the field strength matrix regular at $F = 0$,

$$f_{\text{even}}(F) = \sum_{n=0}^{\infty} c_{2n} F^{2n}, \quad f_{\text{odd}}(F) = \sum_{n=0}^{\infty} c_{2n+1} F^{2n+1} \quad (3.27)$$

Decomposing $\mathcal{G}_{B,F}$ into their even (odd) parts $\mathcal{S}_{B,F}$ ($\mathcal{A}_{B,F}$),

$$\mathcal{G}_{B,F} = \mathcal{S}_{B,F} + \mathcal{A}_{B,F} \quad (3.28)$$

and applying the above formulas we obtain the following matrix decompositions of $\mathcal{G}_B, \dot{\mathcal{G}}_B, \mathcal{G}_F$,

$$\begin{aligned} \mathcal{S}_{B12} &= \frac{T}{2} \left[\frac{A_{B12}^+}{z_+} \hat{\mathcal{Z}}_+^2 + \frac{A_{B12}^-}{z_-} \hat{\mathcal{Z}}_-^2 \right] \\ &= \frac{T}{2(z_+^2 - z_-^2)} \left\{ \left[\frac{z_-^2}{z_+} A_{B12}^+ - \frac{z_+^2}{z_-} A_{B12}^- \right] \mathbb{1} + \left[\frac{A_{B12}^+}{z_+} - \frac{A_{B12}^-}{z_-} \right] \mathcal{Z}^2 \right\} \\ \mathcal{A}_{B12} &= \frac{iT}{2} \left[(S_{B12}^+ - \dot{G}_{B12}) \frac{\hat{\mathcal{Z}}_+}{z_+} + (S_{B12}^- - \dot{G}_{B12}) \frac{\hat{\mathcal{Z}}_-}{z_-} \right] \end{aligned}$$

$$\begin{aligned}
&= \frac{iT}{2(z_+^2 - z_-^2)} \left\{ [S_{B12}^+ - S_{B12}^-] \mathcal{Z} + \left[\frac{z_+}{z_-} (S_{B12}^- - \dot{G}_{B12}) - \frac{z_-}{z_+} (S_{B12}^+ - \dot{G}_{B12}) \right] \tilde{\mathcal{Z}} \right\} \\
\dot{\mathcal{S}}_{B12} &= -S_{B12}^+ \hat{\mathcal{Z}}_+^2 - S_{B12}^- \hat{\mathcal{Z}}_-^2 \\
&= \frac{1}{z_+^2 - z_-^2} \left\{ [z_+^2 S_{B12}^- - z_-^2 S_{B12}^+] \mathbb{1} + [S_{B12}^- - S_{B12}^+] \mathcal{Z}^2 \right\} \\
\dot{\mathcal{A}}_{B12} &= -i [A_{B12}^- \hat{\mathcal{Z}}_- + A_{B12}^+ \hat{\mathcal{Z}}_+] \\
&= \frac{i}{z_+^2 - z_-^2} \left\{ [z_- A_{B12}^- - z_+ A_{B12}^+] \mathcal{Z} + [z_- A_{B12}^+ - z_+ A_{B12}^-] \tilde{\mathcal{Z}} \right\} \\
\mathcal{S}_{F12} &= -S_{F12}^+ \hat{\mathcal{Z}}_+^2 - S_{F12}^- \hat{\mathcal{Z}}_-^2 \\
&= \frac{1}{z_+^2 - z_-^2} \left\{ [z_+^2 S_{F12}^- - z_-^2 S_{F12}^+] \mathbb{1} + [S_{F12}^- - S_{F12}^+] \mathcal{Z}^2 \right\} \\
\mathcal{A}_{F12} &= -i [A_{F12}^- \hat{\mathcal{Z}}_- + A_{F12}^+ \hat{\mathcal{Z}}_+] \\
&= \frac{i}{z_+^2 - z_-^2} \left\{ [z_- A_{F12}^- - z_+ A_{F12}^+] \mathcal{Z} + [z_- A_{F12}^+ - z_+ A_{F12}^-] \tilde{\mathcal{Z}} \right\}
\end{aligned} \tag{3.29}$$

Here we have further introduced

$$z_{\pm} \equiv eN_{\pm}T, \quad \tilde{\mathcal{Z}} \equiv eT\tilde{F}, \quad \mathcal{Z}_{\pm} \equiv eTF_{\pm} = \frac{z_{\pm}^2 \mathcal{Z} - z_+ z_- \tilde{\mathcal{Z}}}{z_{\pm}^2 - z_{\mp}^2}, \quad \hat{\mathcal{Z}}_{\pm} \equiv \frac{\mathcal{Z}_{\pm}}{z_{\pm}} \tag{3.30}$$

Note that $\mathcal{Z}\tilde{\mathcal{Z}} = -z_+ z_- \mathbb{1}$, $\hat{\mathcal{Z}}_{\pm}^3 = -\hat{\mathcal{Z}}_{\pm}$. The scalar, dimensionless coefficient functions appearing in these formulas are given by

$$\begin{aligned}
S_{B12}^{\pm} &= \frac{\sinh(z_{\pm} \dot{G}_{B12})}{\sinh(z_{\pm})} \\
A_{B12}^{\pm} &= \frac{\cosh(z_{\pm} \dot{G}_{B12})}{\sinh(z_{\pm})} - \frac{1}{z_{\pm}} \\
S_{F12}^{\pm} &= G_{F12} \frac{\cosh(z_{\pm} \dot{G}_{B12})}{\cosh(z_{\pm})} \\
A_{F12}^{\pm} &= G_{F12} \frac{\sinh(z_{\pm} \dot{G}_{B12})}{\cosh(z_{\pm})}
\end{aligned} \tag{3.31}$$

The non-vanishing coincidence limits are in $A_{B,F}^\pm$,

$$\begin{aligned} A_{Bii}^\pm &= \coth(z_\pm) - \frac{1}{z_\pm} \\ A_{Fii}^\pm &= \tanh(z_\pm) \end{aligned} \tag{3.32}$$

In the string-inspired formalism, those functions are the basic building blocks of parameter integrals for processes involving constant fields. Let us also write down the first few terms of the weak field expansions of these functions,

$$\begin{aligned} S_{B12}^\pm &= \dot{G}_{B12} \left[1 - \frac{2}{3} \frac{G_{B12}}{T} z_\pm^2 + \left(\frac{2}{45} \frac{G_{B12}}{T} + \frac{2}{15} \frac{G_{B12}^2}{T^2} \right) z_\pm^4 + \mathcal{O}(z_\pm^6) \right] \\ A_{B12}^\pm &= \left(\frac{1}{3} - 2 \frac{G_{B12}}{T} \right) z_\pm + \left(-\frac{1}{45} + \frac{2}{3} \frac{G_{B12}^2}{T^2} \right) z_\pm^3 + \mathcal{O}(z_\pm^5) \\ S_{F12}^\pm &= G_{F12} \left[1 - 2 \frac{G_{B12}}{T} z_\pm^2 + \frac{2}{3} \left(\frac{G_{B12}}{T} + \frac{G_{B12}^2}{T^2} \right) z_\pm^4 + \mathcal{O}(z_\pm^6) \right] \\ A_{F12}^\pm &= G_{F12} \dot{G}_{B12} \left[z_\pm - \left(\frac{1}{3} + \frac{2}{3} \frac{G_{B12}}{T} \right) z_\pm^3 + \mathcal{O}(z_\pm^5) \right] \end{aligned} \tag{3.33}$$

(here we used the identity $\dot{G}_{B12}^2 = 1 - \frac{4}{T} G_{B12}$).

In the same way one finds for the determinant factors (3.11),(3.12)

$$\begin{aligned} \det^{-\frac{1}{2}} \left[\frac{\sin(\mathcal{Z})}{\mathcal{Z}} \right] &= \frac{z_+ z_-}{\sinh(z_+) \sinh(z_-)}, \\ \det^{-\frac{1}{2}} \left[\frac{\tan(\mathcal{Z})}{\mathcal{Z}} \right] &= \frac{z_+ z_-}{\tanh(z_+) \tanh(z_-)} \end{aligned} \tag{3.34}$$

In the appendix we write these formulas also for various special cases of interest. Those are the magnetic field case $\mathbf{E} = 0$, the crossed field case $f = g = 0$, and the self-dual field $F^{\mu\nu} = \tilde{F}^{\mu\nu}$.

Using the above formulas we can obtain explicit results in a Lorentz covariant way. Nevertheless, it will be useful to write down these formulas

also for the Lorentz system where $\mathbf{E}$ and $\mathbf{B}$ are both pointing along the positive z - axis, $\mathbf{E} = (0, 0, E)$, $\mathbf{B} = (0, 0, B)$. (For this to be possible we have to assume that $\mathbf{E} \cdot \mathbf{B} > 0$.) In this Lorentz system $g = iEB$, so that

$$n_{\pm} = \frac{1}{2}(B \pm iE), N_+ = B, N_- = iE, F_+ = Br_{\perp}, F_- = iEr_{\parallel}, \quad (3.35)$$

where

$$r_{\perp} \equiv \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad r_{\parallel} \equiv \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix} \quad (3.36)$$

Introducing also the projectors

$$g_{\perp} \equiv \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad g_{\parallel} \equiv \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (3.37)$$

the matrix decompositions (3.29) can then be rewritten as follows,

$$\begin{aligned}
S_{B12}^{\mu\nu} &= -\frac{T}{2} \sum_{\alpha=\perp,\parallel} \frac{A_{B12}^\alpha}{z_\alpha} g_\alpha^{\mu\nu} \\
\mathcal{A}_{B12}^{\mu\nu} &= \frac{iT}{2} \sum_{\alpha=\perp,\parallel} \frac{S_{B12}^\alpha - \dot{G}_{B12}}{z_\alpha} r_\alpha^{\mu\nu} \\
\dot{S}_{B12}^{\mu\nu} &= \sum_{\alpha=\perp,\parallel} S_{B12}^\alpha g_\alpha^{\mu\nu} \\
\dot{\mathcal{A}}_{B12}^{\mu\nu} &= -i \sum_{\alpha=\perp,\parallel} A_{B12}^\alpha r_\alpha^{\mu\nu} \\
S_{F12}^{\mu\nu} &= \sum_{\alpha=\perp,\parallel} S_{F12}^\alpha g_\alpha^{\mu\nu} \\
\mathcal{A}_{F12}^{\mu\nu} &= -i \sum_{\alpha=\perp,\parallel} A_{F12}^\alpha r_\alpha^{\mu\nu}
\end{aligned} \tag{3.38}$$

with $S/A_{B/F}^\perp \equiv S/A_{B/F}^+(z_+ = eBT \equiv z_\perp)$, $S/A_{B/F}^\parallel \equiv S/A_{B/F}^-(z_- = ieET \equiv z_\parallel)$.

The determinant factors specialize to

$$\begin{aligned}
\det^{-\frac{1}{2}} \left[\frac{\sin(\mathcal{Z})}{\mathcal{Z}} \right] &= \frac{eBT eET}{\sinh(eBT) \sin(eET)} \\
\det^{-\frac{1}{2}} \left[\frac{\tan(\mathcal{Z})}{\mathcal{Z}} \right] &= \frac{eBT eET}{\tanh(eBT) \tan(eET)}
\end{aligned} \tag{3.39}$$

4 The Scalar/Spinor QED Vacuum Polarization Tensors in a Constant Field

We now apply this formalism to a calculation of the scalar and spinor QED vacuum polarisation tensors in a general constant field. For the 2-point case eq.(3.13) yields the following integrand,

$$\exp \left\{ \dots \right\} \big|_{\text{multi-linear}} = \left[\varepsilon_1 \cdot \ddot{\mathcal{G}}_{B12} \cdot \varepsilon_2 - \varepsilon_1 \cdot \dot{\mathcal{G}}_{B1i} \cdot k_i \varepsilon_2 \cdot \dot{\mathcal{G}}_{B2j} \cdot k_j \right] e^{k_1 \cdot \bar{\mathcal{G}}_{B12} \cdot k_2} \tag{4.1}$$

where summation over $i, j = 1, 2$ is understood. Removing the second derivative in the first term by a partial integration in τ_1 this becomes

$$\left[-\varepsilon_1 \cdot \dot{\mathcal{G}}_{B12} \cdot \varepsilon_2 k_1 \cdot \dot{\mathcal{G}}_{B1j} \cdot k_j - \varepsilon_1 \cdot \dot{\mathcal{G}}_{B1i} \cdot k_i \varepsilon_2 \cdot \dot{\mathcal{G}}_{B2j} \cdot k_j \right] e^{k_1 \cdot \bar{\mathcal{G}}_{B12} \cdot k_2} \quad (4.2)$$

We apply the “cycle replacement rule” to this expression and use momentum conservation, $k \equiv k_1 = -k_2$. The content of the brackets then turns into $\varepsilon_{1\mu} I^{\mu\nu} \varepsilon_{2\nu}$, where

$$\begin{aligned} I^{\mu\nu} &= \dot{\mathcal{G}}_{B12}^{\mu\nu} k \cdot \dot{\mathcal{G}}_{B12} \cdot k - \mathcal{G}_{F12}^{\mu\nu} k \cdot \mathcal{G}_{F12} \cdot k \\ &\quad - \left[\left(\dot{\mathcal{G}}_{B11} - \mathcal{G}_{F11} - \dot{\mathcal{G}}_{B12} \right)^{\mu\lambda} \left(\dot{\mathcal{G}}_{B21} - \dot{\mathcal{G}}_{B22} + \mathcal{G}_{F22} \right)^{\nu\kappa} + \mathcal{G}_{F12}^{\mu\lambda} \mathcal{G}_{F21}^{\nu\kappa} \right] k^\kappa k^\lambda \end{aligned} \quad (4.3)$$

Next we would like to use the fact that this integrand contains many terms which integrate to zero due to antisymmetry under the exchange $\tau_1 \leftrightarrow \tau_2$. We therefore decompose $\mathcal{G}_B$ and $\mathcal{G}_F$ into their parts symmetric and antisymmetric in the Lorentz indices as in (3.28). First note that only the even part of $\mathcal{G}_B$ contributes in the exponent,

$$k_1 \cdot \bar{\mathcal{G}}_{B12} \cdot k_2 = k_1 \cdot (\mathcal{S}_{B12} - \mathcal{S}_{B11}) \cdot k_2 \equiv -T k \cdot \Phi_{12} \cdot k \quad (4.4)$$

$I^{\mu\nu}$ turns, after decomposing all factors of $\dot{\mathcal{G}}_B, \mathcal{G}_F$ as above, and deleting all τ - odd terms, into

$$\begin{aligned} I_{\text{spin}}^{\mu\nu} &\equiv \left\{ \left(\dot{\mathcal{S}}_{B12}^{\mu\nu} \dot{\mathcal{S}}_{B12}^{\kappa\lambda} - \dot{\mathcal{S}}_{B12}^{\mu\lambda} \dot{\mathcal{S}}_{B12}^{\nu\kappa} \right) - \left(\mathcal{S}_{F12}^{\mu\nu} \mathcal{S}_{F12}^{\kappa\lambda} - \mathcal{S}_{F12}^{\mu\lambda} \mathcal{S}_{F12}^{\nu\kappa} \right) \right. \\ &\quad \left. + \left(\dot{\mathcal{A}}_{B12} - \dot{\mathcal{A}}_{B11} + \mathcal{A}_{F11} \right)^{\mu\lambda} \left(\dot{\mathcal{A}}_{B12} - \dot{\mathcal{A}}_{B22} + \mathcal{A}_{F22} \right)^{\nu\kappa} \right. \\ &\quad \left. - \mathcal{A}_{F12}^{\mu\lambda} \mathcal{A}_{F12}^{\nu\kappa} \right\} k^\kappa k^\lambda \end{aligned} \quad (4.5)$$

(here we used (3.4)). In this way we obtain the following integral representations for the dimensionally regularised scalar/spinor QED vacuum polarisation tensors [70],

$$\Pi_{\text{scal}}^{\mu\nu}(k) = -\frac{e^2}{[4\pi]^{\frac{D}{2}}} \int_0^\infty \frac{dT}{T} T^{2-\frac{D}{2}} e^{-m^2 T} \det^{-\frac{1}{2}} \left[\frac{\sin(\mathcal{Z})}{\mathcal{Z}} \right] \int_0^1 du_1 I_{\text{scal}}^{\mu\nu} e^{-T k \cdot \Phi_{12} \cdot k} \quad (4.6)$$

$$\Pi_{\text{spin}}^{\mu\nu}(k) = 2 \frac{e^2}{[4\pi]^{\frac{D}{2}}} \int_0^\infty \frac{dT}{T} T^{2-\frac{D}{2}} e^{-m^2 T} \det^{-\frac{1}{2}} \left[\frac{\tan(\mathcal{Z})}{\mathcal{Z}} \right] \int_0^1 du_1 I_{\text{spin}}^{\mu\nu} e^{-T k \cdot \Phi_{12} \cdot k} \quad (4.7)$$

Here $I_{\text{scal}}^{\mu\nu}$ is obtained simply by deleting, in eq. (4.5), all quantities carrying a subscript “F”. As usual we have rescaled to the unit circle and set $u_2 = 0$.

Note that again the transversality of the vacuum polarization tensors is manifest at the integrand level, $k_\mu I_{\text{scal/spin}}^{\mu\nu} = I_{\text{scal/spin}}^{\mu\nu} k_\nu = 0$.

The constant field vacuum polarisation tensors contain the UV divergences of the ordinary vacuum polarisation tensors (2.20),(2.27), and thus require renormalization. As is usual in this context we perform the renormalization on-shell, i.e. we impose the following condition on the renormalized vacuum polarization tensor $\bar{\Pi}^{\mu\nu}(k)$ (see e.g. [5]),

$$\lim_{k^2 \rightarrow 0} \lim_{F \rightarrow 0} \bar{\Pi}^{\mu\nu}(k) = 0 \quad (4.8)$$

Counterterms appropriate to this condition are easy to find from our above results for the ordinary vacuum polarisation tensors. From the representations eqs. (2.18), (2.26) for these tensors it is obvious that we can implement (4.8) by subtracting those same expressions with the last factor $e^{-G_{B12} k^2}$ deleted. In this way we find for the renormalised vacuum polarisation tensors

$$\begin{aligned} \bar{\Pi}_{\text{scal}}^{\mu\nu}(k) &= \Pi_{\text{scal}}^{\mu\nu}(k) + \frac{\alpha}{4\pi} (g^{\mu\nu} k^2 - k^\mu k^\nu) \int_0^\infty \frac{dT}{T} e^{-m^2 T} \int_0^1 du_1 \dot{G}_{B12}^2 \\ \bar{\Pi}_{\text{spin}}^{\mu\nu}(k) &= \Pi_{\text{spin}}^{\mu\nu}(k) - \frac{\alpha}{2\pi} (g^{\mu\nu} k^2 - k^\mu k^\nu) \int_0^\infty \frac{dT}{T} e^{-m^2 T} \int_0^1 du_1 (\dot{G}_{B12}^2 - G_{F12}^2) \end{aligned} \quad (4.9)$$

The remaining u_1 - integral can be brought into a more standard form by a transformation of variables $v = \dot{G}_{B12} = 1 - 2u_1$. Writing the integrands

explicitly using the formulas (3.29) and continuing to Minkowski space ² we obtain our final result for these amplitudes,

$$\begin{aligned}
\bar{\Pi}_{\text{scal}}^{\mu\nu}(k) = & -\frac{\alpha}{4\pi} \int_0^\infty \frac{ds}{s} e^{-ism^2} \int_{-1}^1 \frac{dv}{2} \left\{ \frac{z_+ z_-}{\sinh(z_+) \sinh(z_-)} \right. \\
& \times \exp \left[-i \frac{s}{2} \sum_{\alpha=+,-} \frac{A_{B12}^\alpha - A_{B11}^\alpha}{z_\alpha} k \cdot \hat{\mathcal{Z}}_\alpha^2 \cdot k \right] \\
& \times \sum_{\alpha,\beta=+,-} \left(S_{B12}^\alpha S_{B12}^\beta [(\hat{\mathcal{Z}}_\alpha^2)^{\mu\nu} k \cdot \hat{\mathcal{Z}}_\beta^2 \cdot k - (\hat{\mathcal{Z}}_\alpha^2 k)^\mu (\hat{\mathcal{Z}}_\beta^2 k)^\nu] \right. \\
& \quad \left. - (A_{B12}^\alpha - A_{B11}^\alpha)(A_{B12}^\beta - A_{B22}^\beta)(\hat{\mathcal{Z}}_\alpha k)^\mu (\hat{\mathcal{Z}}_\beta k)^\nu \right) \\
& \left. - (g^{\mu\nu} k^2 - k^\mu k^\nu) v^2 \right\} \tag{4.10}
\end{aligned}$$

$$\begin{aligned}
\bar{\Pi}_{\text{spin}}^{\mu\nu}(k) = & \frac{\alpha}{2\pi} \int_0^\infty \frac{ds}{s} e^{-ism^2} \int_{-1}^1 \frac{dv}{2} \left\{ \frac{z_+ z_-}{\tanh(z_+) \tanh(z_-)} \right. \\
& \times \exp \left[-i \frac{s}{2} \sum_{\alpha=+,-} \frac{A_{B12}^\alpha - A_{B11}^\alpha}{z_\alpha} k \cdot \hat{\mathcal{Z}}_\alpha^2 \cdot k \right] \\
& \times \sum_{\alpha,\beta=+,-} \left([S_{B12}^\alpha S_{B12}^\beta - S_{F12}^\alpha S_{F12}^\beta] [(\hat{\mathcal{Z}}_\alpha^2)^{\mu\nu} k \cdot \hat{\mathcal{Z}}_\beta^2 \cdot k - (\hat{\mathcal{Z}}_\alpha^2 k)^\mu (\hat{\mathcal{Z}}_\beta^2 k)^\nu] \right. \\
& \quad \left. - [(A_{B12}^\alpha - A_{B11}^\alpha + A_{F11}^\alpha)(A_{B12}^\beta - A_{B22}^\beta + A_{F22}^\beta) - A_{F12}^\alpha A_{F12}^\beta] (\hat{\mathcal{Z}}_\alpha k)^\mu (\hat{\mathcal{Z}}_\beta k)^\nu \right) \\
& \left. - (g^{\mu\nu} k^2 - k^\mu k^\nu) (v^2 - 1) \right\} \tag{4.11}
\end{aligned}$$

where now

$$\begin{aligned}
z_+ &= i e s a \\
z_- &= - e s b \\
\hat{\mathcal{Z}}_+ &= \frac{a F - b \tilde{F}}{a^2 + b^2} \\
\hat{\mathcal{Z}}_- &= -i \frac{b F + a \tilde{F}}{a^2 + b^2}
\end{aligned}$$

²For the Maxwell invariants this means $f \rightarrow \mathcal{F}$, $g \rightarrow i\mathcal{G}$, $N_+ \rightarrow a$, $N_- \rightarrow ib$ (to be able to fix all signs we assume $\mathcal{G} \geq 0$). Note also that $r_\perp k \rightarrow \vec{k}_\perp$, $r_\parallel k \rightarrow -i\vec{k}_\parallel$.

$$(4.12)$$

with a, b denoting the standard ‘secular’ invariants

$$\begin{aligned} a &\equiv \sqrt{\sqrt{\mathcal{F}^2 + \mathcal{G}^2} + \mathcal{F}} \\ b &\equiv \sqrt{\sqrt{\mathcal{F}^2 + \mathcal{G}^2} - \mathcal{F}} \end{aligned} \quad (4.13)$$

$$(\mathcal{F} = \tfrac{1}{2}(B^2 - E^2), \quad \mathcal{G} = \mathbf{E} \cdot \mathbf{B}).$$

For fermion QED, the constant field vacuum polarization tensor was obtained before by various authors [20, 21, 22, 23, 24]. For the sake of comparison with their results, let us also specialize to the Lorentz system where $\mathbf{E} = (0, 0, E)$ and $\mathbf{B} = (0, 0, B)$. In this system $a = B, b = E$. Denoting

$$\begin{aligned} k_{\parallel} &= (k^0, 0, 0, k^3), & k_{\perp} &= (0, k^1, k^2, 0) \\ \tilde{k}_{\parallel} &= (k^3, 0, 0, k^0), & \tilde{k}_{\perp} &= (0, k^2, -k^1, 0) \end{aligned} \quad (4.14)$$

our result can be written as follows,

$$\begin{aligned} \bar{\Pi}_{\left(\begin{smallmatrix} \text{spin} \\ \text{scal} \end{smallmatrix}\right)}^{\mu\nu}(k) &= -\frac{\alpha}{4\pi} \begin{pmatrix} -2 \\ 1 \end{pmatrix} \int_0^\infty \frac{ds}{s} \int_{-1}^1 \frac{dv}{2} \left\{ \frac{zz'}{\sin(z) \sinh(z')} \begin{pmatrix} \cos(z) \cosh(z') \\ 1 \end{pmatrix} \right. \\ &\quad \times e^{-is\Phi_0} \sum_{\alpha, \beta = \perp, \parallel} \left[s_{\left(\begin{smallmatrix} \text{spin} \\ \text{scal} \end{smallmatrix}\right)}^{\alpha\beta} (g^{\mu\nu} k_\beta^2 - k_\alpha^\mu k_\beta^\nu) + a_{\left(\begin{smallmatrix} \text{spin} \\ \text{scal} \end{smallmatrix}\right)}^{\alpha\beta} \tilde{k}_\alpha^\mu \tilde{k}_\beta^\nu \right] \\ &\quad \left. - e^{-ism^2} (g^{\mu\nu} k^2 - k^\mu k^\nu) \begin{pmatrix} v^2 - 1 \\ v^2 \end{pmatrix} \right\} \end{aligned} \quad (4.15)$$

where $z = eBs, z' = eEs$,

$$\Phi_0 = m^2 + \frac{k_{\perp}^2}{2} \frac{\cos(zv) - \cos(z)}{z \sin(z)} - \frac{k_{\parallel}^2}{2} \frac{\cosh(z'v) - \cosh(z')}{z' \sinh(z')} \quad (4.16)$$

$$\begin{aligned}
s_{\text{scal}}^{\perp\perp} &= \frac{\sin^2(zv)}{\sin^2(z)} \\
s_{\text{scal}}^{\perp\parallel,\parallel\perp} &= \frac{\sin(zv) \sinh(z'v)}{\sin(z) \sinh(z')} \\
s_{\text{scal}}^{\parallel\parallel} &= \frac{\sinh^2(z'v)}{\sinh^2(z')} \\
a_{\text{scal}}^{\perp\perp} &= \left(\frac{\cos(zv) - \cos(z)}{\sin(z)} \right)^2 \\
a_{\text{scal}}^{\perp\parallel,\parallel\perp} &= -\frac{\cos(zv) - \cos(z)}{\sin(z)} \frac{\cosh(z'v) - \cosh(z')}{\sinh(z')} \\
a_{\text{scal}}^{\parallel\parallel} &= \left(\frac{\cosh(z'v) - \cosh(z')}{\sinh(z')} \right)^2
\end{aligned} \tag{4.17}$$

$$\begin{aligned}
s_{\text{spin}}^{\perp\perp} &= \frac{\sin^2(zv)}{\sin^2(z)} - \frac{\cos^2(zv)}{\cos^2(z)} \\
s_{\text{spin}}^{\perp\parallel,\parallel\perp} &= \frac{\sin(zv) \sinh(z'v)}{\sin(z) \sinh(z')} - \frac{\cos(zv) \cosh(z'v)}{\cos(z) \cosh(z')} \\
s_{\text{spin}}^{\parallel\parallel} &= \frac{\sinh^2(z'v)}{\sinh^2(z')} - \frac{\cosh^2(z'v)}{\cosh^2(z')} \\
a_{\text{spin}}^{\perp\perp} &= \left(\frac{\cos(zv) - \cos(z)}{\sin(z)} - \tan(z) \right)^2 - \frac{\sin^2(zv)}{\cos^2(z)} \\
a_{\text{spin}}^{\perp\parallel,\parallel\perp} &= -\left(\frac{\cos(zv) - \cos(z)}{\sin(z)} - \tan(z) \right) \left(\frac{\cosh(z'v) - \cosh(z')}{\sinh(z')} + \tanh(z') \right) \\
&\quad - \frac{\sin(zv) \sinh(z'v)}{\cos(z) \cosh(z')} \\
a_{\text{spin}}^{\parallel\parallel} &= \left(\frac{\cosh(z'v) - \cosh(z')}{\sinh(z')} + \tanh(z') \right)^2 - \frac{\sinh^2(z'v)}{\cosh^2(z')}
\end{aligned} \tag{4.18}$$

In this form our integrand for the spinor QED case can easily be identified with the one given in [22, 24] using the identities [24]

$$\begin{aligned} g_{\parallel}^{\mu\nu} k_{\parallel}^2 - k_{\parallel}^{\mu} k_{\parallel}^{\nu} &= -\tilde{k}_{\parallel}^{\mu} \tilde{k}_{\parallel}^{\nu} \\ g_{\perp}^{\mu\nu} k_{\perp}^2 - k_{\perp}^{\mu} k_{\perp}^{\nu} &= \tilde{k}_{\perp}^{\mu} \tilde{k}_{\perp}^{\nu} \end{aligned} \quad (4.19)$$

and trigonometric identities³.

For the scalar QED case, the vacuum polarization tensor in a constant field was obtained in [21] using the “operator diagram technique”. The resulting parameter integral representation agrees with our eq.(4.10).

5 Discussion

We have described in detail the application of the ‘string-inspired’ formalism to the calculation of QED photon amplitudes in a constant external field. The two-point cases were studied explicitly. As is usual in the application of the string-inspired technique to QED processes, our calculation of the spinor QED vacuum polarisation tensor in a constant field has yielded the corresponding scalar QED quantity as a byproduct.

In both cases we have arrived at compact results written in terms of the two Lorentz invariants of the Maxwell field. Due to our novel representation (3.29) of the generalized worldline Green’s functions here this was achieved without the use of explicit matrix representations.

Already the two-point case clearly shows the advantages of the method as compared to the standard field theory techniques used in previous calculations of these quantities. There are several aspects to this. The partial integration procedure combined with the Bern-Kosower ‘replacement rule’ effectively replaces the calculation of Dirac traces, and is technically clearly preferable to the latter. Also, the string-inspired technique combines all loop propagators into a single propagator from the beginning, while in a Feynman diagram calculation each propagator has to be parametrized separately, and the global loop proper-time variable is to be introduced at a later stage. The close relationship between the scalar and spinor QED calculations is also

³Up to the global sign, which differs from the one in [22, 24] due to a different definition of the vacuum polarization tensor. (Our field theory conventions follow [71].)

useful. It is due to the fact that the “string-inspired” formalism is based on a second-order field theory formulation of fermion QED [38, 39, 72] rather than the more usual first-order one.

Clearly one would expect these advantages to become more significant with increasing number of legs. For the three-photon amplitude in a magnetic field this was already demonstrated in [46]. We believe that in the present formalism even a calculation of photon-photon scattering in a constant field would not be excessively cumbersome.

In a sequel paper [73] we will consider more generally the calculation of amplitudes involving both vectors and axial vectors in a general constant field, using the generalization of the path integral representation (2.21) given in [74, 75].

Various other generalizations of this formalism would be of interest. An obvious one is the extension to QED amplitudes involving external scalars or fermions; the relevant generalizations of eqs.(2.1),(2.21) have been known for a long time [66, 67, 68]. See [50] (scalar QED) and [76, 77] (spinor QED) for relevant work for the vacuum case. Interesting would also be the extension to the finite temperature case along the lines of [47, 78, 79].

Acknowledgements: The author would like to thank W. Dittrich, L.E. Hernquist, M. Reuter and V.I. Ritus for various helpful informations.

A Worldline Green's functions for special field configurations

In this appendix we give the explicit form of the (Euclidean) generalized worldline Green's functions and determinants for the following three special cases of a constant field, i) the purely magnetic field case, ii) the crossed - field case, iii) the self-dual case.

1. *Magnetic field case:*

$$E = g = 0, f = \frac{1}{2}B^2, n_+ = n_- = \frac{B}{2}, N_- = 0, N_+ = B$$

$$\begin{aligned}
\bar{\mathcal{S}}_{B12} &= G_{B12} \mathbb{1} + \left[\frac{1}{z^2} G_{B12} + T \frac{\cosh(z\dot{G}_{B12}) - \cosh(z)}{2z^3 \sinh(z)} \right] \mathcal{Z}^2 \\
\mathcal{A}_{B12} &= \frac{iT}{2z^2} \left[\frac{\sinh(z\dot{G}_{B12})}{\sinh(z)} - \dot{G}_{B12} \right] \mathcal{Z} \\
\dot{\mathcal{S}}_{B12} &= \dot{G}_{B12} \mathbb{1} + \frac{1}{z^2} \left[\dot{G}_{B12} - \frac{\sinh(z\dot{G}_{B12})}{\sinh(z)} \right] \mathcal{Z}^2 \\
\dot{\mathcal{A}}_{B12} &= i \left[\frac{1}{z^2} - \frac{\cosh(z\dot{G}_{B12})}{z \sinh(z)} \right] \mathcal{Z} \\
\mathcal{S}_{F12} &= G_{F12} \mathbb{1} + G_{F12} \left[\frac{1}{z^2} - \frac{\cosh(z\dot{G}_{B12})}{z^2 \cosh(z)} \right] \mathcal{Z}^2 \\
\mathcal{A}_{F12} &= -iG_{F12} \frac{\sinh(z\dot{G}_{B12})}{z \cosh(z)} \mathcal{Z}
\end{aligned} \tag{A.1}$$

$$\det^{-\frac{1}{2}} \left[\frac{\sin(\mathcal{Z})}{\mathcal{Z}} \right] = \frac{z}{\sinh(z)}, \quad \det^{-\frac{1}{2}} \left[\frac{\tan(\mathcal{Z})}{\mathcal{Z}} \right] = \frac{z}{\tanh(z)} \tag{A.2}$$

where $z = eBT$. We have subtracted from $\mathcal{S}_B$ its coincidence limit, which is expressed by the “bar”. The purely electric case is, of course, analogous.

2. *Crossed field case:*

$$f = g = n_{\pm} = N_{\pm} = 0$$

$$\begin{aligned}
\bar{\mathcal{S}}_{B12} &= G_{B12} \mathbb{1} + \frac{1}{3T} G_{B12}^2 \mathcal{Z}^2 \\
\mathcal{A}_{B12} &= -\frac{i}{3} \dot{G}_{B12} G_{B12} \mathcal{Z} \\
\dot{\mathcal{S}}_{B12} &= \dot{G}_{B12} \mathbb{1} + \frac{2}{3T} \dot{G}_{B12} G_{B12} \mathcal{Z}^2 \\
\dot{\mathcal{A}}_{B12} &= i \left[\frac{2}{T} G_{B12} - \frac{1}{3} \right] \mathcal{Z} \\
\mathcal{S}_{F12} &= G_{F12} \mathbb{1} + \frac{2}{T} G_{F12} G_{B12} \mathcal{Z}^2 \\
\mathcal{A}_{F12} &= -i G_{F12} \dot{G}_{B12} \mathcal{Z}
\end{aligned} \tag{A.3}$$

$$\det^{-\frac{1}{2}} \left[\frac{\sin(\mathcal{Z})}{\mathcal{Z}} \right] = \det^{-\frac{1}{2}} \left[\frac{\tan(\mathcal{Z})}{\mathcal{Z}} \right] = 1 \tag{A.4}$$

3. *Self dual case:*

$$F = \tilde{F}, f = g, n_- = 0, n_+ = N_+ = N_- = \sqrt{f}$$

$$\begin{aligned}
\bar{\mathcal{S}}_{B12} &= \frac{T \cosh(Z) - \cosh(Z \dot{G}_{B12})}{2 Z \sinh(Z)} \mathbb{1} \\
\mathcal{A}_{B12} &= i \frac{T}{2Z^2} \left[\frac{\sinh(Z \dot{G}_{B12})}{\sinh(Z)} - \dot{G}_{B12} \right] \mathcal{Z} \\
\dot{\mathcal{S}}_{B12} &= \frac{\sinh(Z \dot{G}_{B12})}{\sinh(Z)} \mathbb{1} \\
\dot{\mathcal{A}}_{B12} &= -i \left[\frac{\cosh(Z \dot{G}_{B12})}{Z \sinh(Z)} - \frac{1}{Z^2} \right] \mathcal{Z} \\
\mathcal{S}_{F12} &= G_{F12} \frac{\cosh(Z \dot{G}_{B12})}{\cosh(Z)} \mathbb{1}
\end{aligned}$$

$$\mathcal{A}_{F12} = -iG_{F12} \frac{\sinh(Z\dot{G}_{B12})}{Z \cosh(Z)} \mathcal{Z} \quad (\text{A.5})$$

$$\det^{-\frac{1}{2}} \left[\frac{\sin(\mathcal{Z})}{\mathcal{Z}} \right] = \frac{Z^2}{\sinh^2(Z)}, \quad \det^{-\frac{1}{2}} \left[\frac{\tan(\mathcal{Z})}{\mathcal{Z}} \right] = \frac{Z^2}{\tanh^2(Z)} \quad (\text{A.6})$$

where $Z = \sqrt{f}eT$.

References

- [1] H. Euler and B. Kockel, *Naturwissenschaften* **23** (1935) 246.
- [2] W. Heisenberg and H. Euler, *Z. Phys.* **98** (1936) 714.
- [3] V. Weisskopf, K. Dan. Vidensk. Selsk. Mat. Fy. Medd. **14** (1936) 1, reprinted in *Quantum Electrodynamics*, J. Schwinger (Ed.) (Dover, New York, 1958).
- [4] V. I. Ritus, *The Lagrangian Function of an Intense Electromagnetic Field and Quantum Electrodynamics at Short Distances*, in *Proc. Lebedev Phys. Inst. vol. 168*, V. I. Ginzburg ed., Nova Science Publ., NY 1987.
- [5] W. Dittrich and M. Reuter, *Effective Lagrangians in Quantum Electrodynamics*, (Springer, 1985).
- [6] W. Greiner and J. Reinhardt, *Quantum Electrodynamics* (Springer, 1992).
- [7] W. Dittrich and H. Gies, “Vacuum Birefringence in Strong Magnetic Fields”, *Proc. of Frontier Tests of Quantum Electrodynamics and Physics of the Vacuum*, Sandansky 1998, 1998 Heron Press, 29, (hep-ph/9806417).
- [8] R. Pengo et al., Workshop Sandansky 1998, loc. cit., 59.
- [9] V. Fock, *Physik. Z. Sowjetunion* **12** (1937) 404.
- [10] J. Schwinger, *Phys. Rev.* **82** (1951) 664.
- [11] J. Toll, PhD thesis, Princeton University 1952 (unpublished).
- [12] A. Minguzzi, *Nuovo Cim.* **6** (1957) 501; **9** (1958) 145; **19** (1961) 847.
- [13] R. Baier and P. Breitenlohner, *Acta Phys. Austr.* **25** (1967) 212; *Nuovo Cim.* **47** (1967) 261.
- [14] D.H. Constantinescu, *Nucl. Phys.* **B 36** (1972) 121.
- [15] W.-Y. Tsai and T. Erber, *Phys. Rev.* **D 10** (1974) 492.

- [16] R.A. Cover and G. Kalman, Phys. Rev. Lett. **33** (1974) 1113.
- [17] A.E. Shabad, Ann. Phys. **90** (1975) 166.
- [18] N.B. Narozhnyi, Sov. Phys. JETP **28** (1969) 371.
- [19] V.I. Ritus, Ann. Phys. **69** (1972) 555.
- [20] I.A. Batalin and A.E. Shabad, Sov. Phys. JETP **33** (1971) 483.
- [21] V. N. Baier, V. M. Katkov, and V. M. Strakhovenko, Sov. Phys. JETP **40** (1975) 225; Sov. Phys. JETP **41** (1975) 198.
- [22] L.F. Urrutia, Phys. Rev. **D 17** (1978) 1977.
- [23] G.K. Artimovich, Sov. Phys. JETP **70** (1990) 787.
- [24] H. Gies, PhD thesis, Tübingen University (1999); hep-ph/9909500.
- [25] Z. Bialynicka-Birula and I. Bialynicka-Birula, Phys. Rev. **D 2** (1970) 2341.
- [26] S.L. Adler, Ann. Phys. NY **67** (1971) 599.
- [27] G.G. Raffelt, *Stars as Laboratories for Fundamental Physics*, University of Chicago Press, Chicago 1996.
- [28] M. G. Baring and A. K. Harding (astro-ph/9704210).
- [29] V.M. Kaspi et al., to appear in the Proc. of *Stellar Endpoints, AGN and the Diffuse Background*, Bologna 1999.
- [30] D. Bakalov, INFN/AE-94/27 (unpublished).
- [31] S.A. Lee et al., “Measurement of the Magnetically-induced QED Birefringence of the Vacuum and an Improved Laboratory Search for Light Pseudoscalars”, FERMILAB-PROPOSAL-P-877A, 1998.
- [32] J. Géhéniau, Physica **16** (1950) 822.
- [33] W. Y. Tsai, Phys. Rev. **D 10** (1974) 1342.

- [34] Z. Bern, D.A. Kosower, Phys. Rev. **D 38** (1988) 1888; Phys. Rev. Lett. **66** (1991) 1669; Nucl. Phys. **B379** (1992) 451.
- [35] Z. Bern, L. Dixon, D. A. Kosower, Phys. Rev. Lett. **70** (1993) 2677 (hep-ph/9302280).
- [36] Z. Bern, D. C. Dunbar, and T. Shimada, Phys. Lett. **B312**(1993) 277 (hep-th/9307001).
- [37] D.C. Dunbar and P. S. Norridge, Nucl. Phys. **B 433** (1995) 181 (hep-th/9408014).
- [38] Z. Bern and D. C. Dunbar, Nucl. Phys. **B379** (1992) 562.
- [39] M. J. Strassler, Nucl. Phys. **B385** (1992) 145.
- [40] M.G. Schmidt and C. Schubert, Phys. Rev. **D53** (1996) 2150 (hep-th/9410100).
- [41] M. G. Schmidt and C. Schubert, Phys. Lett. **B318** (1993) 438 (hep-th/9309055).
- [42] D. Cangemi, E. D'Hoker, and G. Dunne, Phys. Rev. **D 51** (1995) 2513.
- [43] V.P. Gusynin and I.A. Shovkovy, J. Phys. **74** (1996) 282 (hep-ph/9509383); UCTP-106-98 (hep-th/9804143).
- [44] R. Shaisultanov, Phys. Lett. **B 378** (1996) 354 (hep-th/9512142).
- [45] M. Reuter, M.G. Schmidt, and C. Schubert, Ann. Phys. **259**, 313 (1997) (hep-th/9610191).
- [46] S. L. Adler and C. Schubert, Phys. Rev. Lett. **77**, 1695 (1996) (hep-th/9605035).
- [47] I.A. Shovkovy, Phys. Lett. **B441** (1998) 313 (hep-th/9806156).
- [48] D. Fliegner, M.G. Schmidt, and C. Schubert, Z. Phys. **C 64** (1994) 111 (hep-ph/9401221); and with P. Haberl, Ann. Phys. (N.Y.) **264** (1998) 51 (hep-th/9707189).

- [49] M. G. Schmidt and C. Schubert, Phys. Lett. **B331** (1994) 69 (hep-th/9403158).
- [50] K. Daikouji, M. Shino, and Y. Sumino, Phys. Rev. **D53** (1996) 4598 (hep-ph/9508377).
- [51] K. Roland and H.-T. Sato, Nucl. Phys. **B480** 99 (1996) (hep-th/9604152).
- [52] D. Fliegner, M. Reuter, M.G. Schmidt, and C. Schubert, Theor. Math. Phys. **113** (1997) 1442 (hep-th/9704194).
- [53] B. Körs and M.G. Schmidt, Eur. Phys. J. **C6** (1999) 175 (hep-th/9803144).
- [54] G. V. Dunne and C. Schubert, Nucl. Phys. **B 564** (2000) 591 (hep-th/9907190).
- [55] Z. Bern, TASI Lectures, Boulder TASI 92, 471 (hep-ph/9304249).
- [56] C. Schubert, Lectures given at 36th Cracow School of Theoretical Physics, Zakopane 1996, Act. Phys. Polon. **B 27** (1996) 3965 (hep-th/9610108).
- [57] H.M. Fried, *Functional Methods and Models in Quantum Field Theory*, MIT Press, Cambridge (1972).
- [58] A. Barducci, F. Bordi, and R. Casalbuoni, Nuovo Cim. **64 B** (1981) 287.
- [59] I.A. Batalin, E.S. Fradkin, and Sh.M. Shvartsman, Nucl. Phys. **B258** (1985) 435.
- [60] S.G. Rajeev, Ann. Phys. **173** (1987) 249.
- [61] E.S. Fradkin, D.M. Gitman, and S.M. Shvartsman, *Quantum Electrodynamics with Unstable Vacuum*, Springer 1991.
- [62] D.G.C. McKeon and T.N. Sherry, Mod. Phys. Lett. **A9** (1994) 2167.
- [63] R. P. Feynman, Phys. Rev. **80** (1950) 440.

- [64] M.B. Green, J.H. Schwarz, and E. Witten, *Superstring Theory*, Cambridge University Press 1987.
- [65] C. Itzykson and J. Zuber, *Quantum Field Theory*, McGraw-Hill 1985.
- [66] E. S. Fradkin, Nucl. Phys. **76** (1966) 588.
- [67] R. Casalbuoni, Nuov. Cim. **33A** (1976) 389.
- [68] F.A. Berezin and M.S. Marinov, Ann. Phys. **104** (1977) 336.
- [69] C. Schubert, Eur. Phys. J. **C5** (1998) 693 (hep-th/9710067).
- [70] C. Schubert, LAPTH-Conf-729/99, to appear in the Proc. of *Corfu Summer Institute on Elementary Particle Physics, 1998* (hep-ph/9905525).
- [71] S. Weinberg, *The Quantum Theory of Fields*, Vol. 1, Cambridge Univ. Pr. 1995.
- [72] A. Morgan, Phys. Lett. **B351** (1995) 249 (hep-ph/9502230).
- [73] C. Schubert, in preparation.
- [74] D.G.C. McKeon and C. Schubert, Phys. Lett. **B440** (1998) 101 (hep-th/9807072).
- [75] F.A. Dilkes, D.G.C. McKeon, and C. Schubert, J. High Energy Phys. **03** (1999) 022 (hep-th/9812213).
- [76] D.G.C. McKeon and A. Rebhan, Phys. Rev. **D48** (1993) 2891.
- [77] A.I. Karanikas and C.N. Ktorides, JHEP 9911:033 (hep-th/9905027).
- [78] M. Haack and M.G. Schmidt, Eur. Phys. J. **C7** (1999) 149 (hep-th/9806138).
- [79] H.-T. Sato, HD-THEP-98-41 (hep-th/9809053).